

Taxonomic study of the genus *Ischnothyreus* Simon, 1893 from Myanmar (Araneae, Oonopidae)

Yanfeng Tong^{1,2}, Shuqiang Li³, Dongju Bian⁴

1 Life Science College, Shenyang Normal University, Shenyang 110034, China **2** Southeast Asia Biological Diversity Research Institute, Chinese Academy of Sciences, Yezin, Nay Pyi Taw 05282, Myanmar **3** Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China **4** CAS Key Laboratory of Forest Ecology and Management, Institute of Applied Ecology, Shenyang 110016, China

Corresponding author: Shuqiang Li (lisq@ioz.ac.cn); Dongju Bian (biandongju@163.com)

Academic editor: I. Agnarsson | Received 17 August 2020 | Accepted 19 October 2020 | Published 16 November 2020

<http://zoobank.org/DAFA5BEE-7571-4CA8-99D1-D76DFFCAB875>

Citation: Tong Y, Li S, Bian D (2020) Taxonomic study of the genus *Ischnothyreus* Simon, 1893 from Myanmar (Araneae, Oonopidae). ZooKeys 993: 1–26. <https://doi.org/10.3897/zookeys.993.57676>

Abstract

Seven new species of the genus *Ischnothyreus* Simon, 1893 from the spider family Oonopidae Simon, 1890 are reported from Myanmar: *I. hponkanrazi* **sp. nov.** (♀), *I. jianglangi* **sp. nov.** (♀), *I. meukyawwa* **sp. nov.** (♂♀), *I. putao* **sp. nov.** (♀), *I. qiuxing* **sp. nov.** (♀), *I. taunggyi* **sp. nov.** (♂♀) and *I. zhigangi* **sp. nov.** (♂♀). Morphological descriptions and photographic illustrations of the new species are given. All types are preserved in the Institute of Zoology, Chinese Academy of Sciences in Beijing (IZCAS).

Keywords

Goblin spider, morphology, new species, taxonomy

Introduction

The genus *Ischnothyreus* was established by Simon in 1893, with *Ischnaspis peltifer* Simon, 1892 from Saint Vincent in the Caribbean as the type species (Simon 1893). This genus is an Old World taxon, being represented in the New World by only two species, *I. peltifer* and *I. velox* Jackson, 1908, both of which are assumed to be introduced (Platnick et al. 2012; Brescovit et al. 2019).

The genus *Ischnothyreus* Simon, 1893 can be recognized by the presence of leg spines, the usually small abdominal scutum, the strongly sclerotized male palps, the heavily sclerotized male endites, and the winding genital tube in females (Kranz-Baltensperger 2011). There are currently 107 valid specific names assigned to *Ischnothyreus* (Li 2020; WSC 2020) and even more are waiting to be described (Richard et al. 2016).

In this paper seven new *Ischnothyreus* species collected from Myanmar are described and illustrated. This work presents the first record and description of species *Ischnothyreus* from Myanmar.

Materials and methods

The specimens were examined in 95% ethanol using a Leica M205C stereomicroscope. Details were studied with an Olympus BX51 compound microscope. Photos were taken with a Canon EOS 750D zoom digital camera (18 megapixels) mounted on an Olympus BX51 compound microscope. Vulvae were cleared in lactic acid. Scanning electron microscope images (SEM) were taken under high vacuum with a Hitachi TM3030 after critical point drying and gold-palladium coating. All measurements were taken using an Olympus BX51 compound microscope and are given in millimeters in the text. The specimens are preserved in the Institute of Zoology, Chinese Academy of Sciences (IZCAS) in Beijing, China (curator: Jun Chen).

The following abbreviations are used in the text and figures: **a** = apodemes; **ALE** = anterior lateral eyes; **ass** = anchor-shaped structure; **bsa** = bell-shaped atrium; **csa** = circular atrium; **csp** = crown-shaped sclerotized process; **hsm** = hook-shaped membrane; **llm** = leaf like membrane; **lpp** = leaf-shaped prolateral projection; **nlm** = needle like membrane; **nsa** = nipple-shaped atrium; **oa** = opening of the atrium; **PLE** = posterior lateral eyes; **PME** = posterior median eyes; **rl** = retrolateral lobe; **sem** = serrated exterior margin; **tsa** = triangular shaped atrium; **vpr** = ventral protuberance; **wt** = winding tube.

Taxonomy

Family Oonopidae Simon, 1890

Genus *Ischnothyreus* Simon, 1893

Ischnothyreus hponkanrazi Tong & Li, sp. nov.

<http://zoobank.org/EFA6509C-6C65-46CC-B65D-E54770EAB9E5>

Figures 1, 16A, B

Type material. *Holotype* ♀: MYANMAR, Kachin State, Putao, Hponkanrazi Wildlife Sanctuary, roadside between Camp 2 to Camp 1; 27°36'067"N, 96°59'367"E; elevation ca 1714 m; 17.XII.2016; Wu J. leg. (IZCAS AR-25158).




Figure 1. *Ischnothyreus hponkanrazi* sp. nov., female holotype **A–C** habitus, dorsal, ventral and lateral views **D–G** prosoma, dorsal, ventral, lateral and anterior views **H** epigastric region, ventral view. Scale bars: 0.4 mm (**A–F**); 0.2 mm (**G, H**).

Diagnosis. The new species is similar to *I. campanaceus* Tong & Li, 2008 in the bell-shaped atrium, but can be distinguished by the short abdominal dorsal scutum (1/3 of the abdomen length (Fig. 1A) vs 4/5 of the abdomen length (Tong and Li 2008: fig. 1B; Tong 2013: fig. 44B), and the greater sinuosity of the winding tube of endogyne (Fig. 1H) (vs short, simple winding tube; Tong and Li 2008: fig. 1F; Tong 2013: fig. 44F).

Description. Female (holotype). Body: habitus as in Fig. 1A–C; body length 2.40. **Carapace:** 1.08 long, 0.98 wide; pale brown, without any pattern, ovoid in dorsal view, slightly elevated in lateral view, surface finely reticulate, fovea absent, lateral margin straight, smooth (Fig. 1D, F). **Clypeus:** height about equal to ALE radius or more (Fig. 1G). **Eyes:** six, in one group, well developed, subequal, ALE circular, PME and PLE oval, posterior eye row procurved from both above and front (Fig. 1D, G). **Sternum:** as long as wide, pale orange, uniform, not fused to carapace, surface smooth, setae sparse (Fig. 1E). **Mouthparts:** chelicerae, endites and labium orange; chelicerae straight, base of fangs unmodified (Fig. 1G); labium rectangular, not fused to sternum, anterior margin not indented at middle; endites unmodified (Fig. 1E). **Abdomen:** 1.33 long, 0.74 wide; dorsal scutum well sclerotized, pale brown, covering 1/3 of the abdomen width and approximately 1/3 of the abdomen length, not fused to epigastric scutum; epigastric and postgastric scutum well sclerotized, pale orange, unfused (Fig. 1A, B). **Legs:** pale orange, femur I with two prolateral spines, tibia I with four pairs, metatarsus I with two pairs of long ventral spines. Leg II spination is similar to leg I except femur with only one prolateral spine. Legs III and IV spineless. **Epigastric area:** surface without external features (Fig. 1H). **Endogyne:** from the middle of the slightly thickened margin of the postgastric scutum runs a dark winding tube posteriorly, ending in a bell-shaped atrium (Fig. 16A, B).

Male. Unknown.

Etymology. The specific name is a noun in apposition taken from the type locality.

Distribution. Known only from the type locality.

Ischnothyreus jianglangi Tong & Li, sp. nov.

<http://zoobank.org/505F6C65-291C-47C0-8997-23CA63EA5161>

Figures 2, 17A, B

Type material. Holotype ♀: MYANMAR, Kachin State, Putao, Hponkanrazi Wildlife Sanctuary, roadside between Camp 2 to Camp 1; 27°35'806"N, 96°59'532"E; elevation ca 1613 m; 10.V.2017; Wu J. & Chen Z. leg. (IZCAS AR-25159).

Diagnosis. The new species is similar to *I. jojo* Kranz-Baltensperger, 2011 in the triangular shaped atrium, but can be distinguished by shape of the dorsal abdominal scutum (width/length = 1/2 (Fig. 2A) vs approximately 1/3; Kranz-Baltensperger 2011: fig. 25A), and the unmodified exterior surface of postgastric scutum (Fig. 17A) (vs with curved, sclerotized extensions and U-shaped structure; Kranz-Baltensperger 2011: fig. 25E, F).




Figure 2. *Ischnothyreus jianglangi* sp. nov., female holotype **A–C** habitus, dorsal, ventral and lateral views **D–G** prosoma, dorsal, ventral, lateral and anterior views **H** epigastric region, ventral view. Scale bars: 0.4 mm (**A–F**); 0.1 mm (**G, H**).

Description. Female (holotype). *Body:* habitus as in Fig. 2A–C; body length 2.35. *Carapace:* 1.02 long, 0.87 wide; brown, without any pattern, ovoid in dorsal view, slightly elevated in lateral view, surface finely reticulate, fovea absent, lateral margin straight, smooth (Fig. 2D, F). *Clypeus:* height about equal to ALE radius or less (Fig. 2G). *Eyes:* six, in one group, well developed, subequal, ALE circular, PME and PLE oval, posterior eye row procurved from both above and front (Fig. 2D, G). *Sternum:* as long as wide, pale brown, uniform, not fused to carapace, surface smooth, setae sparse (Fig. 2E). *Mouthparts:* chelicerae, endites and labium pale brown; chelicerae straight, base of fangs unmodified (Fig. 2G); labium rectangular, not fused to sternum, anterior margin not indented at middle; endites unmodified (Fig. 2E). *Abdomen:* 1.42 long, 1.01 wide; dorsal scutum covering less than 1/2 of the abdomen length and 1/3 of the abdomen width, not fused to epigastric scutum; epigastric and postgastric scutum well sclerotized, pale orange, unfused (Fig. 2A, B). *Legs:* pale orange, femur I with two prolateral spines, tibia I with four pairs, metatarsus I with two pairs of long ventral spines. Leg II spination is similar to leg I except femur with only one prolateral spine. Legs III and IV spineless. *Epigastric area:* surface without external features (Fig. 2H). *Endogyne:* from the middle of the slightly thickened margin of the postgastric scutum runs a dark, simple winding tube posteriorly, ending in a black, triangular shaped atrium (Fig. 17A, B).

Male. Unknown.

Etymology. The species is named after Mr Jianglang Wu, one of the collectors of the holotype; noun in genitive case.

Distribution. Known only from the type locality.

Ischnothyreus meukyawwa Tong & Li, sp. nov.

<http://zoobank.org/08A2A68D-453F-4E7F-8F52-05FB0C49EDAC>

Figures 3–5, 14A–C, 15A, B, 16C, D

Type material. Holotype ♂: MYANMAR, Kachin State, Putao, Meukyawwa Village; 27°20'883"N, 97°22'717"E; elevation ca 464 m; 25.XII.2016; Wu J. leg. (IZCAS AR-25160). **Paratypes:** 5♂ 9♀, same data as for holotype (IZCAS AR-25161–25174).

Diagnosis. The new species is similar to *I. an* Tong & Li, 2016 in the large abdominal dorsal and ventral scutum, but can be distinguished by the unmodified male cheliceral fang (Fig. 3H, I) (vs with thorn-like protrusion; Tong et al. 2016: fig. 1G, H), the acute distal end of male palp (Fig. 14A) (vs blunt distal end of male palp; Tong et al. 2016: fig. 3A, D), and the small bell-shaped atrium (Fig. 16C) (vs a large equilateral triangular shaped atrium; Tong et al. 2016: fig. 2G, I).

Description. Male (holotype). *Body:* habitus as in Fig. 3A–C; body length 1.57. *Carapace:* 0.89 long, 0.74 wide; pale brown, with egg-shaped patches behind eyes, ovoid in dorsal view, strongly elevated in lateral view, surface of elevated portion of pars cephalica smooth, sides finely reticulate, fovea absent, lateral margin straight, smooth




Figure 3. *Ischnothyreus meukyawwa* sp. nov., male holotype **A–C** habitus, dorsal, ventral and lateral views **D–G** prosoma, dorsal, ventral, lateral and anterior views **H, I** left chelicerae, anterior and posterior views. Scale bars: 0.4 mm (**A–F**); 0.2 mm (**G**); 0.1 mm (**H, I**).




Figure 4. *Ischnothyreus meukyauwa* sp. nov., male holotype, left palp, SEM **A, B, G** prolateral, retrolateral and dorsal views **C, D** palp bulb, prolateral and retrolateral views **E, F, H** distal part of palpal bulb, prolateral, retrolateral and dorsal views. Abbreviations: hsm = hook-shaped membrane; lpp = leaf-shaped prolateral projection; rl = retrolateral lobe; vpr = ventral protuberance.




Figure 5. *Ischnothyreus meukyawwa* sp. nov., female paratype **A–C** habitus, dorsal, ventral and lateral views **D** prosoma, anterior view **E** epigastric region, ventral view. Abbreviation: **ass** = anchor-shaped structure. Scale bars: 0.4 mm (**A–C**); 0.2 mm (**D, E**).

(Fig. 3D, F). **Clypeus:** height about $2/3$ of ALE diameter (Fig. 3G). **Eyes:** six, in one group, well developed, subequal, ALE circular, PME and PLE oval, posterior eye row recurved from above, procurved from front (Fig. 3D, G). **Sternum:** as long as wide, pale orange, uniform, not fused to carapace, surface smooth, setae sparse (Fig. 3E). **Mouthparts:** chelicerae, endites and labium orange; chelicerae straight, base of fangs with crown-shaped sclerotized process with serrated exterior margin, fang groove with a few small denticles (Figs 3H, I, 15A, B); labium rectangular, not fused to sternum, anterior margin not indented at middle; anteromedian tip of endites with one strong, tooth-like projection (Fig. 3E). **Abdomen:** 0.84 long, 0.43 wide; dorsal scutum well sclerotized, pale orange, covering whole abdomen width and approximately $5/6$ of the

abdomen length, not fused to epigastric scutum; epigastric and postgastric scutum well sclerotized, pale orange, fused, postgastric scutum covering about 5/6 of the abdomen length (Fig. 3A–C). **Legs:** pale orange, femur I with three prolateral and two small retrolateral spines, tibia I with four pairs, metatarsus I with two pairs of long ventral spines. Leg II spination is similar to leg I except femur with only two prolateral and one retrolateral spine. Legs III and IV spineless. **Palp:** strongly sclerotized, trochanter with ventral projection, cymbium brown, fused with bulb; bulb brown, with one large and one very small ventral protuberance, distal end of bulb elongated, with one broad leaf-shaped prolateral projection and distal hook-shaped membrane, retrolateral lobe narrow (Figs 4, 14A–C).

Female (paratype, IZCAS AR-25160). Same as male except as noted. **Body:** habitus as in Fig. 5A–C; body length 1.93. **Carapace:** 0.89 long, 0.76 wide; without any pattern. **Mouthparts:** chelicerae and endites unmodified. **Abdomen:** 0.83 long, 0.75 wide; dorsal scutum covering 3/5 of the abdomen length, about 1/2 of the abdomen width. **Epigastric area:** the postgastric scutum with central anchor-shaped structure (Fig. 5E). **Endogyne:** from the middle of the slightly thickened margin of the postgastric scutum runs a dark, simple winding tube posteriorly, ending in a small bell-shaped atrium (Fig. 16C, D).

Etymology. The specific name is a noun in apposition taken from the type locality.

Distribution. Known only from the type locality.

Ischnothyreus putao Tong & Li, sp. nov.

<http://zoobank.org/D96DFBF0-A942-44A3-833A-18EB8FB325B1>

Figures 6, 17C, D

Type material. Holotype ♀: MYANMAR, Kachin State, Putao, Hponkanrazi Wildlife Sanctuary, roadside between Camp 1 to Camp 2; 27°36'567"N, 96°58'850"E; elevation ca 2233 m; 15.XII.2016; Wu J. leg. (IZCAS AR-25175).

Diagnosis. The new species is similar to *I. zhigangi* sp. nov. in the very small abdominal dorsal scutum and the brown carapace, but can be distinguished by the large bell-shaped atrium (Fig. 17C) (vs small, bell-shaped atrium; Fig. 17E), and the smoothly curved posterior margin of postgastric scutum (Fig. 6H) (vs straight posterior margin; Fig. 13H).

Description. Female (holotype). **Body:** habitus as in Fig. 6A–C; body length 1.94. **Carapace:** 0.97 long, 0.88 wide; dark brown, without any pattern, ovoid in dorsal view, strongly elevated in lateral view, surface finely reticulate, fovea absent, lateral margin straight, smooth (Fig. 6D, F). **Clypeus:** height about equal to ALE radius or more (Fig. 6G). **Eyes:** six, in one group, well developed, subequal, ALE circular, PME and PLE oval, posterior eye row procurved from both above and front (Fig. 6D, G). **Sternum:** as long as wide, pale brown, uniform, not fused to carapace, surface smooth, setae sparse (Fig. 6E). **Mouthparts:** chelicerae, endites and labium brown; chelicerae straight, base of fangs unmodified (Fig. 6G); labium rectangular, not fused to sternum, anterior margin not indented at middle; endites unmodified (Fig. 6E). **Abdomen:** 1.30




Figure 6. *Ischnothyreus putao* sp. nov., female holotype **A–C** habitus, dorsal, ventral and lateral views **D–G** prosoma, dorsal, ventral, lateral and anterior views **H** epigastric region, ventral view. Abbreviation: ass = anchor-shaped structure. Scale bars: 0.4 mm (**A–F**); 0.2 mm (**G**); 0.1 mm (**H**).

long, 0.89 wide; dorsal scutum weakly sclerotized, very small, not fused to epigastric scutum; epigastric and postgastric scutum well sclerotized, pale orange, unfused (Fig. 6A, B). **Legs:** pale orange, femur I with three prolateral spines, tibia I with four pairs, metatarsus I with two pairs of long ventral spines. Leg II spination is similar to leg I except femur with only two prolateral spine. Legs III and IV spineless. **Epigastric area:** the postgastric scutum with central, anchor-shaped structure, and smoothly curved posterior margin (Fig. 6H). **Endogyne:** from the middle of the slightly thickened margin of the postgastric scutum runs a dark, simple winding tube posteriorly, ending in a large, inverted bell-shaped atrium (Fig. 17C, D).

Male. Unknown.

Etymology. The specific name is a noun in apposition taken from the type locality.

Distribution. Known only from the type locality.

Ischnothyreus qiuxing Tong & Li, sp. nov.

<http://zoobank.org/8B8870A1-6AD1-4C30-B8E3-616837C148B7>

Figures 7, 16E, F

Type material. Holotype ♀: MYANMAR, Kachin State, Putao, Around Ziradum Village; 27°33'465"N, 97°06'580"E; elevation ca 1051 m; 8.V.2017; Wu J. & Chen Z. leg. (IZCAS AR-25176).

Diagnosis. The new species is similar to *I. balu* Kranz-Baltensperger, 2011 in the circular atrium, but can be distinguished by the size of atrium (nearly 1/5 the length of postgastric scutum (Fig. 16E) vs more than 1/3 the length of postgastric scutum; Kranz-Baltensperger 2011: fig. 2F, H) and the greater sinuosity of the winding tube (Fig. 16F) (vs short, simple winding tube; Kranz-Baltensperger 2011: fig. 2G).

Description. Female (holotype). Body: habitus as in Fig. 7A–C; body length 2.01. **Carapace:** 0.87 long, 0.72 wide; yellow, without any pattern, ovoid in dorsal view, strongly elevated in lateral view, surface finely reticulate, fovea absent, lateral margin straight, smooth (Fig. 7D, F). **Clypeus:** height about equal to ALE radius or less (Fig. 7G). **Eyes:** six, in one group, well developed, subequal, ALE circular, PME and PLE oval, posterior eye row recurved from above, procurved from front (Fig. 7D, G). **Sternum:** as long as wide, yellow, uniform, not fused to carapace, surface smooth, setae sparse (Fig. 7E). **Mouthparts:** chelicerae, endites and labium orange; chelicerae straight, base of fangs unmodified (Fig. 7G); labium rectangular, not fused to sternum, anterior margin not indented at middle; endites unmodified (Fig. 7E). **Abdomen:** 1.35 long, 0.92 wide; dorsal scutum weakly sclerotized, very small, not fused to epigastric scutum; epigastric and postgastric scutum well sclerotized, pale orange, unfused (Fig. 7A, B). **Legs:** pale orange, femur I with two prolateral spines, tibia I with four pairs, metatarsus I with two pairs of long ventral spines. Leg II spination is similar to leg I except femur with only one prolateral spine. Legs III and IV spineless. **Epigastric area:** surface without external features (Fig. 7H). **Endogyne:** from the middle of the




Figure 7. *Ischnothyreus qiuxing* sp. nov., female holotype **A–C** habitus, dorsal, ventral and lateral views **D–G** prosoma, dorsal, ventral, lateral and anterior views **H** epigastric region, ventral view. Scale bars: 0.4 mm (**A–C**); 0.2 mm (**D–H**).

slightly thickened margin of the postgastric scutum runs a dark, very simple winding tube posteriorly, ending in a circular atrium (Fig. 16E, F).

Male. Unknown.

Etymology. The specific name is derived from Chinese pinyin, “qixing”, which means “circular”, referring to the circular atrium; noun in apposition.

Distribution. Known only from the type locality.

***Ischnothyreus taunggyi* Tong & Li, sp. nov.**

<http://zoobank.org/E8228172-15A8-4BCB-8B68-BC87903BCBDE>

Figures 8–10, 14D–F, 15C, D, 16G, H

Type material. Holotype ♂: MYANMAR, Shan State, Taunggyi, East of Nyaung Shwe Township; 20°34'700"N, 96°57'450"E; elevation ca 1005 m; 30.XI.2016; Wu J. leg. (IZCAS AR-25177). **Paratypes** 2♀: same data as for holotype (IZCAS AR-25178–25179).

Diagnosis. The new species is similar to *I. zhigangi* sp. nov. in the male palp and the crown-shaped sclerotized process of male cheliceral fang, but can be distinguished by the long abdominal dorsal scutum (3/4 of the abdomen length (Fig. 8A) vs very small; Fig. 11A) and ventral scutum (4/5 of the abdomen length (Fig. 8B) vs very small; Fig. 11B) of male, and the long abdominal dorsal scutum (less than 1/2 of the abdomen length (Fig. 10A) vs very small; Fig. 13A) and the nipple-shaped atrium (Fig. 16G) (vs inverted bell-shaped atrium; Fig. 17E) of female.

Description. Male (holotype). Body: habitus as in Fig. 8A–C; body length 1.71. **Carapace:** 0.84 long, 0.63 wide; pale brown, with egg-shaped patches behind eyes, ovoid in dorsal view, strongly elevated in lateral view, surface of elevated portion of pars cephalica smooth, sides finely reticulate, fovea absent, lateral margin straight, smooth (Fig. 8D). **Clypeus:** height about equal to ALE radius or more. **Eyes:** six, in one group, well developed, ALE largest, ALE circular, PME and PLE oval, posterior eye row recurved from above, procurved from front (Fig. 8D). **Sternum:** as long as wide, pale orange, uniform, not fused to carapace, surface smooth, setae sparse (Fig. 8E). **Mouthparts:** chelicerae, endites and labium orange; chelicerae straight, with crown-shaped sclerotized process at base of fangs, fang groove with a few small and one larger denticles (Fig. 15C, D); labium rectangular, not fused to sternum, anterior margin not indented at middle; anteromedian tip of endites with one strong, tooth-like projection (Fig. 8E). **Abdomen:** 0.74 long, 0.49 wide; dorsal scutum well sclerotized, pale orange, covering 2/3 the abdomen width and approximately 3/4 of the abdomen length, unfused to epigastric scutum; epigastric and postgastric scutum well sclerotized, pale orange, fused, postgastric scutum covering about 4/5 of the abdomen length (Fig. 8A, B). **Legs:** pale orange, femur I with three prolateral and one small retrolateral spines, tibia I with four pairs, metatarsus I with two pairs of long ventral spines. Leg II spination is similar to leg I except femur with only two prolateral spines. Legs III and IV spineless. **Palp:** strongly sclerotized, trochanter with ventral projection, cymbium brown, fused with bulb; bulb brown, with two large ventral protuberances, distal end of bulb




Figure 8. *Ischnothyreus taunggyi* sp. nov., male holotype **A–C** habitus, dorsal, ventral and lateral views **D–F** prosoma, dorsal, ventral and lateral views **G, H** left chelicerae, anterior and posterior views. Scale bars: 0.4 mm (**A–C**); 0.2 mm (**D–F**); 0.1 mm (**G, H**).




Figure 9. *Ischnothyreus taunggyi* sp. nov., male holotype, left palp, SEM **A, B** prolateral and retrolateral views **C, D, G** palpal bulb, prolateral, retrolateral and dorsal views **E, F, H** distal part of palpal bulb, prolateral, retrolateral and dorsal views. Abbreviations: nlm = needle like membrane; llm = leaf like membrane; lpp = leaf-shaped prolateral projection; rl = retrolateral lobe; vpr = ventral protuberance.




Figure 10. *Ischnothyreus taunggyi* sp. nov., female paratype **A–C** habitus, dorsal, ventral and lateral views **D–F, G** prosoma, dorsal, ventral, lateral and anterior views **H** epigastric region, ventral view. Abbreviation: ass = anchor-shaped structure. Scale bars: 0.4 mm (**A–F**); 0.2 mm (**G, H**).

elongated, with one leaf-shaped prolateral projection and distal needle like membrane, retrolateral lobe broad, ear-shaped (Figs 9, 14D–F).

Female (paratype, IZCAS AR-25178). Same as male except as noted. **Body:** habitus as in Fig. 10A–C; body length 1.98. **Carapace:** 0.83 long, 0.71 wide; without any pattern (Fig. 10D). **Mouthparts:** chelicerae and endites unmodified (Fig. 10E, G). **Abdomen:** 1.27 long, 0.83 wide; dorsal scutum covering less than 1/2 of the abdomen length, about 1/3 of the abdomen width (Fig. 10A). **Epigastric area:** the postgastric scutum with central anchor-shaped structure (Fig. 10H). **Endogyne:** from the middle of the slightly thickened margin of the postgastric scutum runs a dark, very complex winding tube posteriorly, ending in a small, bell-shaped atrium (Fig. 16G, H).

Etymology. The specific name is a noun in apposition taken from the type locality.

Distribution. Known only from the type locality.

Ischnothyreus zhigangi Tong & Li, sp. nov.

<http://zoobank.org/CB6C29F2-AC7C-4511-8F20-7771D82CB329>

Figures 11–13, 14G–I, 15E, F, 17E, F

Type material. **Holotype** ♂: MYANMAR, Putao, Hponkanrazi Wildlife Sanctuary Around Camp 2; 27°36'681"N, 96°58'958"E; elevation ca 2457 m; 11.V.2017; Wu J. & Chen Z. leg. (IZCAS AR-25178). **Paratypes** 4♀: data same as for holotype; 27°31'103"N, 96°57'694"E; elevation ca 2737 m; 16.V.2017; Wu J. & Chen Z. leg. (IZCAS AR-25179–25182).

Diagnosis. The new species is similar to *I. taunggyi* sp. nov. but can be distinguished by the short abdominal dorsal scutum (very small (Fig. 11A) vs 3/4 of the abdomen length; Fig. 8A) and ventral scutum (very small (Fig. 11B) vs 4/5 of the abdomen length; Fig. 8B) of male, and the short abdominal dorsal scutum (very small (Fig. 13A) vs less than 1/2 of the abdomen length; Fig. 10A) and the inverted bell-shaped atrium (Fig. 17E) (vs nipple-shaped atrium (Fig. 16G) of female).

Description. Male (holotype). **Body:** habitus as in Fig. 11A–C; body length 2.08. **Carapace:** 1.03 long, 0.87 wide; pale brown, with egg-shaped patches behind eyes, ovoid in dorsal view, strongly elevated in lateral view, surface of elevated portion of pars cephalica smooth, sides finely reticulate, fovea absent, lateral margin straight, smooth (Fig. 11D). **Clypeus:** height about equal to ALE radius or more. **Eyes:** six, in one group, well developed, subequal, ALE circular, PME and PLE oval, posterior eye row straight from above, procurved from front (Fig. 11D). **Sternum:** as long as wide, pale brown, uniform, not fused to carapace, surface smooth, setae sparse (Fig. 11E). **Mouthparts:** chelicerae, endites and labium orange; chelicerae straight, base of fangs with crown-shaped sclerotized process with serrated exterior margin (Figs 11G, H, 15E, F), fang groove with a few small denticles; labium rectangular, not fused to sternum, anterior margin not indented at middle; anteromedian tip of endites with one strong, tooth-like projection (Fig. 11E). **Abdomen:** 1.07 long, 0.72 wide; dorsal scutum weakly sclerotized, pale orange, very small, not fused to epigastric scutum; epigastric and




Figure 11. *Ischnothyreus zbigangi* sp. nov., male holotype **A–C** habitus, dorsal, ventral and lateral views **D–F** prosoma, dorsal, ventral and lateral views **G, H** left chelicerae, anterior and posterior views. Scale bars: 0.4 mm (**A–F**); 0.1 mm (**G, H**).




Figure 12. *Ischnothyreus zbigangi* sp. nov., male holotype, left palp, SEM **A, B** prolateral and retrolateral views **C, D, G** palpal bulb, prolateral, retrolateral and dorsal views **E, F, H** distal part of palpal bulb, prolateral, retrolateral and dorsal views. Abbreviations: lpp = leaf-shaped prolateral projection; rl = retrolateral lobe; vpr = ventral protuberance.




Figure 13. *Ischnothyreus zhigangi* sp. nov., female paratype **A–C** habitus, dorsal, ventral and lateral views **D–G** prosoma, dorsal, ventral, lateral and anterior views **H** epigastric region, ventral view. Abbreviation: ass = anchor-shaped structure. Scale bars: 0.4 mm (**A–G**); 0.1 mm (**H**).



Figure 14. *Ischnothyreus* spp., left male palp **A–C** *I. meukyawwa* sp. nov. **D–F** *I. taunggyi* sp. nov. **G–I** *I. zbigangi* sp. nov. **A, D, G** prolateral view **B, E, H** dorsal view **C, F, I** retrolateral view. Scale bars: 0.1 mm.




Figure 15. *Ischnothyreus* spp., left male chelicerae. **A, B** *I. meukyauwa* sp. nov. **C, D** *I. taunggyi* sp. nov. **E, F** *I. zbigangi* sp. nov. **A, C, E** anterior view **B, D, F** posterior view. Abbreviations: csp = crown-shaped sclerotized process; sem = serrated exterior margin.

postgastric scutum weakly sclerotized, pale orange, fused, postgastric scutum very small (Fig. 11A, B). **Legs:** pale orange, femur I with three prolateral spines, tibia I with four pairs, metatarsus I with two pairs of long ventral spines. Leg II spination is similar to leg I except femur with only two prolateral spines. Legs III and IV spineless. **Palp:** strongly sclerotized, trochanter with ventral projection, cymbium brown, fused with bulb; bulb brown, with one large ventral protuberance, distal end of bulb elongated, with one narrow leaf-shaped prolateral projection, retrolateral lobe small, simple (Figs 12, 14G–H).




Figure 16. *Ischnothyreus* spp., female copulatory organ **A, B** *I. hponkanrazi* sp. nov. **C, D** *I. meukyawwa* sp. nov. **E, F** *I. qiuxing* sp. nov. **G, H** *I. taunggyi* sp. nov. **A, C, E, G** ventral view **B, D, F, H** dorsal view. Abbreviations: a = apodemes; ass = anchor-shaped structure; bsa = bell-shaped atrium; csa = circular atrium; nsa = nipple-shaped atrium; oa = opening of the atrium; wt = winding tube. Scale bars: 0.2 mm.

Female (paratype, IZCAS AR-25179). Same as male except as noted. **Body:** habitus as in Fig. 13A–C; body length 2.63. **Carapace:** 1.10 long, 0.95 wide; without any pattern, posterior eye row procurved from both above and front (Fig. 13D, G). **Mouthparts:** chelicerae and endites unmodified (Fig. 13E, G). **Abdomen:** 1.82 long, 1.19 wide. **Epigastric area:** the postgastric scutum with central anchor-shaped structure (Fig. 13H). **Endogyne:** from the middle of the slightly thickened margin of the




Figure 17. *Ischnothyreus* spp., female copulatory organ **A, B** *I. jianglangi* sp. nov. **C, D** *I. putao* sp. nov. **E, F** *I. zbigangi* sp. nov. **A, C, E** ventral view **B, D, F** dorsal view. Abbreviations: a = apodemes; ass = anchor-shaped structure; bsa = bell-shaped atrium; tsa = triangular shaped atrium; wt = winding tube. Scale bars: 0.1 mm

postgastric scutum runs a dark, simple winding tube posteriorly, ending in a small, inverted bell-shaped atrium (Fig. 17E, F).

Etymology. The species is named after Mr Zhigang Chen, one of the collectors of the holotype; noun in genitive case.

Distribution. Known only from the type locality.

Acknowledgements

The manuscript benefited greatly from constructive comments by Ingi Agnarsson, Antonio Domingos Brescovit, Lily Berniker, and Nadine Dupérré. This study was supported by the National Natural Science Foundation of China (31750002, 31972867) and the Program for Liaoning Innovation Talents in University to Yanfeng Tong, and by the National Natural Science Foundation of China (NSFC-31530067), and the Southeast Asia Biodiversity Research Institute, Chinese Academy of Sciences (2015CASEABRI005, Y4ZK111B01) to Shuqiang Li.

References

- Brescovit AD, Bonaldo AB, Ott R, Chavari JL (2019) To boldly go: on invasive goblin spiders in Brazil (Araneae, Oonopidae). *Iheringia, Série Zoologia* 109: e2019033. <https://doi.org/10.1590/1678-4766e2019033>
- Kranz-Baltensperger Y (2011) The oonopid spider genus *Ischnothyreus* in Borneo (Oonopidae, Araneae). *Zootaxa* 2939: 1–49. <https://doi.org/10.11646/zootaxa.2939.1.1>
- Li S (2020) Spider taxonomy for an advanced China. *Zoological Systematics* 45(2): 73–77.
- Platnick NI, Berniker L, Kranz-Baltensperger Y (2012) The goblin spider genus *Ischnothyreus* (Araneae, Oonopidae) in the New World. *American Museum Novitates* 3759: 1–32. <https://doi.org/10.1206/3759.2>
- Richard M, Graber W, Kropf C (2016) The goblin spider genus *Ischnothyreus* (Araneae, Oonopidae) in Java and Sumatra. *Zootaxa* 4151(1): 1–99. <https://doi.org/10.11646/zootaxa.4151.1.1>
- Simon E (1893) *Histoire naturelle des araignées*. Deuxième édition, tome premier. Roret, Paris, 257–488. <https://doi.org/10.5962/bhl.title.51973>
- Tong Y, Li S (2008) The oonopid spiders (Araneae: Oonopidae) from Hainan Island, China. *Raffles Bulletin of Zoology* 56: 55–66.
- Tong Y (2013) *Haplogynae Spiders from Hainan, China*. Science Press, Beijing, 96 pp. [81 pls]
- Tong Y, Koh JKH, Tong X, Li S (2016) Five new species of the genus *Ischnothyreus* Simon, 1893 from Singapore. *ZooKeys* 618: 39–66. <https://doi.org/10.3897/zookeys.618.9451>
- WSC (2020) *World Spider Catalog, version 21.5*. Natural History Museum Bern. <http://wsc.nmbe.ch> [Accessed on: 2020.08.12]